

Short communication

Detection of antibodies to *Neospora caninum* in two species of wild canids, *Lycalopex gymnocercus* and *Cercdocyon thous* from Brazil

W.A. Cañón-Franco^a, L.E.O. Yai^b, S.L.P. Souza^a, L.C. Santos^c,
N.A.R. Farias^d, J. Ruas^d, F.W. Rossi^e, A.A.B. Gomes^f,
J.P. Dubey^g, S.M. Gennari^{a,*}

^a Departamento de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, Avenida Prof. Orlando Marques de Paiva, 87, Cidade Universitária São Paulo, São Paulo, CEP 05508-000, Brazil

^b Centro de Controle de Zoonoses da Prefeitura do Município de São Paulo, São Paulo, Brazil

^c Itaipu Binacional, Foz do Iguaçu, Paraná, Brazil

^d Instituto de Biologia, Universidade Federal de Pelotas, Rio Grande de Sul, Brazil

^e Divisão Técnica de Medicina Veterinária e Manejo da Fauna Silvestre da Prefeitura do Município de São Paulo, São Paulo, Brazil

^f Departamento de Medicina Veterinária Preventiva, Centro de Saúde e Tecnologia Rural, Universidade Federal da Paraíba, Brazil

^g Animal Parasitic Diseases Laboratory, United States Department of Agriculture, Agriculture Research Service, Animal and Natural Resources Institute, Building 1001, Beltsville, MD 20705, USA

Accepted 4 June 2004

Abstract

Domestic dog (*Canis domesticus*) and the coyote (*Canis latrans*) are the only known definitive hosts for the protozoan *Neospora caninum* that causes abortion in dairy cattle. In the present study, antibodies to *N. caninum* were sought in three species of wild canids, *Cercdocyon thous*, *Lycalopex gymnocercus* and *Dusicyon vetulus* from Brazil. Antibodies to *N. caninum* were assayed by the indirect fluorescent antibody test (IFAT) and the *Neospora* agglutination test (NAT). *N. caninum* antibodies were found in five of 12 *L. gymnocercus* with IFAT titers of 1:50 in three, 1:100 in one, and 1:1600 in one, and NAT titers of 1:40, 1:80, 1:160, 1:320, and 1:640 in five animals. Antibodies to *N. caninum* were found in four of 15 *C. thous* with IFAT titers of 1:50 in one, and

* Corresponding author.

E-mail address: sgennari@usp.br (S.M. Gennari).

1:100 in three, and NAT titer of 1:40 in one animal. All 30 *D. ventulus* were seronegative by IFAT and NAT.

© 2004 Elsevier B.V. All rights reserved.

Keywords: *Neospora caninum*; Wild canids; *Cerdocyon thous*; *Lycalopex gymnocercus*; *Dusicyon vetulus*; Antibodies; Brazil

1. Introduction

The coccidium *Neospora caninum* is an important cause of abortion in cattle worldwide (Dubey and Lindsay, 1996; Dubey, 2003a). It also causes mortality in sheep, goats, horses, rhinoceros, and deer. Domestic dog (*Canis domesticus*) and the coyote (*Canis latrans*) are the two known definitive hosts for *N. caninum* that excrete the environmentally resistant oocysts (McAllister et al., 1998; Basso et al., 2001; Gondim et al., 2004). Because both dogs and coyotes excrete only a few *N. caninum* oocysts compared with other coccidians, other wild carnivores have not been ruled out as its definitive hosts. Antibodies to *N. caninum* have been reported in several species of domestic and wild animals (Dubey, 2003b). We report antibodies to *N. caninum* for the first time in two species of wild canids from Brazil.

2. Materials and methods

Sera were collected from three species of wild canids (*Lycalopex gymnocercus*, *Dusicyon vetulus*, and *Cerdocyon thous*) captured in traps and were part of larger project on the epidemiology and control of infectious diseases in wild animals in Brazil. The *L. gymnocercus* ($n = 12$) were from the state of Rio Grande do Sul, *D. vetulus* ($n = 30$) were from the state of Paraíba, and *C. thous* ($n = 15$) from the state of São Paulo and Paraná.

Blood samples were collected from a jugular or brachial vein, the sera were separated, and stored at -20°C until being tested for anti-*N. caninum* antibodies. The indirect immunofluorescent antibody test (IFAT) with a cut-off value of 1:50 and *Neospora* agglutination test (NAT) with cut-off value of 1:40 were used to screen for *N. caninum* antibodies. The seropositive samples were tested further in two-fold serial dilutions.

For IFAT, tachyzoites of the *N. caninum* NC-1 strain were used as antigen (Dubey et al., 1988). Anti-dog IgG–FITC conjugate from rabbit (Sigma, F7884, St. Louis, Missouri) was used as the fluorescent label, and a positive and a negative domestic dog sera were used in each slide as controls. The NAT was performed using mouse-derived *N. caninum* tachyzoites as described by Romand et al. (1998).

3. Results and discussion

N. caninum antibodies were found in five of 12 *L. gymnocercus* with IFAT titers of 1:50 in three, 1:100 in one, and 1:1600 in one, and NAT titers of 1:40, 1:80, 1:160, 1:320, and 1:640 in five animals. Antibodies to *N. caninum* were found in four of 15 *C. thous* with

IFAT titers of 1:50 in one, and 1:100 in three, and NAT titer of 1:40 in one animal. All 30 *D. vetulus* were seronegative by IFAT and NAT. Results of the present study indicate that additional species of carnivores have been exposed to *N. caninum*. It remains to be determined if these animals can shed *N. caninum* oocysts.

References

- Basso, W., Venturini, L., Venturini, M.C., Hill, D.E., Kwok, O.C.H., Shen, S.K., Dubey, J.P., 2001. First isolation of *Neospora caninum* from the feces of a naturally infected dog. *J. Parasitol.* 87, 612–618.
- Dubey, J.P., 2003a. Neosporosis in cattle. *J. Parasitol.* 89 (Suppl.), S42–S56.
- Dubey, J.P., 2003b. Review of *Neospora caninum* and neosporosis in animals. *Korean J. Parasitol.* 41, 1–16.
- Dubey, J.P., Lindsay, D.S., 1996. A review of *Neospora caninum* and neosporosis. *Vet. Parasitol.* 67, 1–59.
- Dubey, J.P., Hattel, A.L., Lindsay, D.S., Topper, M.J., 1988. Neonatal *Neospora caninum* infection in dogs: isolation of the causative agent and experimental transmission. *J. Am. Vet. Med. Assoc.* 193, 1259–1263.
- Gondim, L.F.P., McAllister, M.M., Pitt, W.C., Zemlicka, D.E., 2004. Coyotes (*Canis latrans*) are definitive hosts of *Neospora caninum*. *Int. J. Parasitol.* 34, 159–161.
- McAllister, M.M., Dubey, J.P., Lindsay, D.S., Jolley, W.R., Wills, R.A., McGuire, A.M., 1998. Dogs are definitive hosts of *Neospora caninum*. *Int. J. Parasitol.* 28, 1473–1478.
- Romand, S., Thulliez, P., Dubey, J.P., 1998. Direct agglutination test for serologic diagnosis of *Neospora caninum* infection. *Parasitol. Res.* 84, 50–53.